

Approximation properties of Chlodowsky variant of (p, q) Bernstein-Stancu-Schurer operators

Vishnu Narayan Mishra^{a,b}, M. Mursaleen^{c,1}, Shikha Pandey^a

^aDepartment of Applied Mathematics & Humanities, Sardar Vallabhbhai National Institute of Technology, Ichchhanath Mahadev Dumas Road, Surat -395 007 (Gujarat), India

^bL. 1627 Awadh Puri Colony Beniganj, Phase -III, Opp. Industrial Training Institute, Ayodhya Main Road, Faizabad-224 001, (Uttar Pradesh), India

^cDepartment of Mathematics, Aligarh Muslim University, Aligarh-202002, India

Abstract

In the present paper, we introduce the Chlodowsky variant of (p, q) Bernstein-Stancu-Schurer operators which is a generalization of (p, q) Bernstein-Stancu-Schurer operators. We have also discussed its Korovkin type approximation properties and rate of convergence.

Keywords: (p, q) -integers; (p, q) -Bernstein operators; (p, q) Bernstein-Stancu-Schurer operators; modulus of continuity; linear positive operators; Korovkin type approximation theorem.

2000 Mathematics Subject Classification: Primary 41A25, 41A36, 41A10, 41A30.

1. Introduction and preliminaries

In 1912, S.N. Bernstein [1] introduced the following sequence of operators $B_n : C[0, 1] \rightarrow C[0, 1]$ defined for any $n \in \mathbb{N}$ and for any function $f \in C[0, 1]$,

$$B_n(f; x) = \sum_{k=0}^n \binom{n}{k} x^k (1-x)^{n-k} f\left(\frac{k}{n}\right), \quad x \in [0, 1]. \quad (1.1)$$

Later various generalizations of these operators were discovered. It has been proved as a powerful tool for numerical analysis, computer aided geometric design and solutions of differential equations. In last two decades, the applications of q -calculus has played an important role in the area of approximation theory, number theory and theoretical physics. In 1987, Lupaş [6] and in 1997, Phillips [16] introduced a sequence of Bernstein polynomials based on q -integers and investigated its approximation properties. Several researchers obtained various other generalizations of operators based on q -calculus. For any function $f \in C[0, 1]$ the q -form of Bernstein operator is described by Lupaş [6] as

$$L_{n,q}(f; x) = \sum_{k=0}^n \frac{\begin{bmatrix} n \\ k \end{bmatrix}_q q^{k(k-1)/2} x^k (1-x)^{n-k}}{\prod_{j=0}^{n-1} (1-x+q^j x)} f\left(\frac{[k]_q}{[n]_q}\right) \quad x \in [0, 1]. \quad (1.2)$$

Email addresses: vishnunarayanmishra@gmail.com, vishnu_narayanmishra@yahoo.co.in (Vishnu Narayan Mishra), mursaleenm@gmail.com (M. Mursaleen), sp1486@gmail.com (Shikha Pandey)

¹Corresponding author

In 1932, Chlodowsky [3] presented a generalization of Bernstein polynomials on an unbounded set, known as Bernstein-Chlodowsky polynomials given by,

$$B_n(f, x) = \sum_{k=0}^n f\left(\frac{k}{n}b_n\right) \binom{n}{k} \left(\frac{x}{b_n}\right)^k \left(1 - \frac{x}{b_n}\right)^{n-k}, \quad 0 \leq x \leq b_n, \quad (1.3)$$

where b_n is an increasing sequence of positive terms with the properties $b_n \rightarrow \infty$ and $\frac{b_n}{n} \rightarrow 0$ as $n \rightarrow \infty$. In 2008, Karsli and Gupta [5] expressed the q -analogue of Bernstein-Chlodowsky polynomials by

$$C_n(f; q; x) = \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_q \left(\frac{x}{b_n}\right)^k \prod_{s=0}^{n-k-1} (1 - q^s \frac{x}{b_n}) f\left(\frac{[k]_q}{[n]_q} b_n\right), \quad 0 \leq x \leq b_n, \quad (1.4)$$

where b_n is an increasing sequence of positive values, with the properties $b_n \rightarrow \infty$ and $\frac{b_n}{[n]_q} \rightarrow 0$ as $n \rightarrow \infty$.

Recently, Mursaleen et al. [7, 9, 11] proposed and analysed approximation properties for (p, q) analogue of Bernstein operators, Bernstein-Stancu operators and Bernstein-Schurer operators. Besides this, we also refer to some recent related work on this topic: e.g. ([12], [13], [14], [15], [17]). Now, we recall certain notations of (p, q) -calculus.

For $0 < q < p < 1$, the (p, q) integer $[n]_{p,q}$ is described by

$$[n]_{p,q} := \frac{p^n - q^n}{p - q}.$$

(p, q) factorial is expressed as

$$[n]_{p,q}! = [n]_{p,q} [n-1]_{p,q} [n-2]_{p,q} \dots 1.$$

(p, q) binomial coefficient is expressed as

$$\left[\begin{matrix} n \\ k \end{matrix} \right]_{p,q} := \frac{[n]_{p,q}!}{[k]_{p,q}! [n-k]_{p,q}!}.$$

(p, q) binomial expansion as

$$(ax + by)_{p,q}^n := \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_{p,q} a^{n-k} b^k x^{n-k} y^k.$$

$$(x + y)_{p,q}^n := (x + y)(px + qy)(p^2x + q^2y) \dots (p^{n-1}x + q^{n-1}y).$$

In 2015, Mursleen et al [10], investigated (p, q) form of Bernstein-Stancu operator, which is given by

$$S_n^{(\alpha, \beta)}(f; x, p, q) = \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^n \left[\begin{matrix} n \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} x^k \prod_{s=0}^{n+m-k-1} (p^s - q^s x) f\left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta}\right), \quad (1.5)$$

where α, β are non-negative integers and $f \in C[0, 1]$, $x \in [0, 1]$ and $0 \leq \alpha \leq \beta$.

For the first few moments, we have the following lemma:

Lemma 1. (See [10]) For the Operators $S_n^{(\alpha, \beta)}$, we have

1. $S_n^{(\alpha, \beta)}(1; x, p, q) = 1,$
2. $S_n^{(\alpha, \beta)}(t; x, p, q) = \frac{[n]_{p,q}x + \alpha}{[n]_{p,q} + \beta},$
3. $S_n^{(\alpha, \beta)}(t^2; x, p, q) = \frac{1}{([n]_{p,q} + \beta)^2} (q[n]_{p,q}[n-1]_{p,q}x^2 + [n]_{p,q}(2\alpha + p^{n-1})x + \alpha^2).$

2. Construction of the operators

Considering the revised form of (p, q) analogue of Bernstein operators (see [8]), we construct the Chlodowsky variant of (p, q) Bernstein-Stancu-Schurer operators as

$$C_{n,m}^{(\alpha, \beta)}(f; x, p, q) = \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^{k} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) f \left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n \right), \quad (2.1)$$

where $n \in \mathbb{N}$, $m, \alpha, \beta \in \mathbb{N}_0$ with $0 \leq \alpha \leq \beta$, $0 \leq x \leq b_n$, $0 < q < p < 1$. Evidently, $C_{n,m}^{(\alpha, \beta)}$ is a linear and positive operator. Consider the case, if $p, q \rightarrow 1$ and $m = 0$ in (2.1), then it will reduce to the Stancu-Chlodowsky polynomials [4].

To begin with, we obtain the following lemma, which will be used throughout the paper.

Lemma 2. Let $C_{n,m}^{(\alpha, \beta)}(f; x, p, q)$ be given by (2.1). The first few moments of the operators are

- (i) $C_{n,m}^{(\alpha, \beta)}(1; x, p, q) = 1,$
- (ii) $C_{n,m}^{(\alpha, \beta)}(t; x, p, q) = \frac{[n+m]_{p,q}x + \alpha b_n}{[n]_{p,q} + \beta},$
- (iii) $C_{n,m}^{(\alpha, \beta)}(t^2; x, p, q) = \frac{1}{([n]_{p,q} + \beta)^2} (q[n+m]_{p,q}[n+m-1]_{p,q}x^2 + [n+m]_{p,q}(2\alpha + p^{n-1})b_nx + \alpha^2b_n^2),$
- (iv) $C_{n,m}^{(\alpha, \beta)}((t-x); x, p, q) = \left(\frac{[n+m]_{p,q}}{[n]_{p,q} + \beta} - 1 \right) x + \frac{\alpha b_n}{[n]_{p,q} + \beta},$
- (v)

$$\begin{aligned} C_{n,m}^{(\alpha, \beta)}((t-x)^2; x, p, q) &= \left(1 - 2 \frac{[n+m]_{p,q}}{([n]_{p,q} + \beta)} + \frac{q[n+m]_{p,q}[n+m-1]_{p,q}}{([n]_{p,q} + \beta)^2} \right) x^2 \\ &\quad + \left(\frac{(2\alpha + p^{n-1})[n+m]_{p,q}}{([n]_{p,q} + \beta)} - 2\alpha \right) \frac{b_n}{([n]_{p,q} + \beta)} x + \frac{\alpha^2 b_n^2}{([n]_{p,q} + \beta)^2}. \end{aligned}$$

Proof. (i)

$$C_{n,m}^{(\alpha, \beta)}(1; x, p, q) = \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) = 1,$$

(ii)

$$\begin{aligned}
C_{n,m}^{(\alpha,\beta)}(t; x, p, q) &= \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n \right) \\
&= \frac{[n+m]_{p,q}}{p^{\frac{n(n-3)}{2}} ([n]_{p,q} + \beta)} \sum_{k=0}^{n+m-1} \left[\begin{matrix} n+m-1 \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k+1)}{2}} \left(\frac{x}{b_n} \right)^{k+1} \prod_{s=0}^{n+m-k-2} (p^s - q^s \frac{x}{b_n}) \left(\frac{b_n}{p^{k+1}} \right) \\
&\quad + \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{\alpha}{[n]_{p,q} + \beta} b_n \right) \\
&= \frac{[n+m]_{p,q}}{([n]_{p,q} + \beta)} x \frac{1}{p^{\frac{(n-1)(n-2)}{2}}} \sum_{k=0}^{n+m-1} \left[\begin{matrix} n+m-1 \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-2} (p^s - q^s \frac{x}{b_n}) + \frac{\alpha b_n}{([n]_{p,q} + \beta)} \\
&= \frac{[n+m]_{p,q}}{([n]_{p,q} + \beta)} x + \frac{\alpha b_n}{([n]_{p,q} + \beta)},
\end{aligned}$$

(iii)

$$\begin{aligned}
C_{n,m}^{(\alpha,\beta)}(t^2; x, p, q) &= \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n \right)^2 \\
&= \frac{1}{p^{\frac{n(n-1)}{2}}} \frac{1}{([n]_{p,q} + \beta)^2} \left[p^{2n} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \frac{b_n^2 [k]_{p,q}^2}{p^{2k}} \right. \\
&\quad + 2\alpha p^n \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \frac{b_n^2 [k]_{p,q}}{p^k} \\
&\quad \left. + \alpha^2 \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) b_n^2 \right] \\
&= \frac{1}{([n]_{p,q} + \beta)^2} \left[\frac{p^{2n}}{p^{\frac{n(n-1)}{2}}} [n+m]_{p,q} b_n^2 \sum_{k=0}^{n+m-1} \left[\begin{matrix} n+m-1 \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k+1)}{2}} \left(\frac{x}{b_n} \right)^{k+1} \prod_{s=0}^{n+m-k-2} (p^s - q^s \frac{x}{b_n}) \frac{[k+1]_{p,q}}{p^{2(k+1)}} \right. \\
&\quad \left. + \frac{p^n}{p^{\frac{n(n-1)}{2}}} 2\alpha [n+m]_{p,q} b_n^2 \sum_{k=0}^{n+m-1} \left[\begin{matrix} n+m-1 \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k+1)}{2}} \left(\frac{x}{b_n} \right)^{k+1} \prod_{s=0}^{n+m-k-2} (p^s - q^s \frac{x}{b_n}) \frac{1}{p^{(k+1)}} + \alpha^2 b_n^2 \right].
\end{aligned}$$

Now using $[k+1]_{p,q} = p^k + q[k]_{p,q}$, we will achieve the result.

Using linear property of operators, we have

$$\begin{aligned}
C_{n,m}^{(\alpha,\beta)}((t-x); x, p, q) &= C_{n,m}^{(\alpha,\beta)}(t; x, p, q) - x C_{n,m}^{(\alpha,\beta)}(1; x, p, q) \\
&= \left(\frac{[n+m]_{p,q}}{[n]_{p,q} + \beta} - 1 \right) x + \frac{\alpha b_n}{[n]_{p,q} + \beta}.
\end{aligned}$$

Hence, we get (iv).

Similar calculations give

$$C_{n,m}^{(\alpha,\beta)}((t-x)^2; x, p, q) = C_{n,m}^{(\alpha,\beta)}(t^2; x, p, q) - 2xC_{n,m}^{(\alpha,\beta)}(t; x, p, q) + x^2C_{n,m}^{(\alpha,\beta)}(1; x, p, q).$$

Substituting the results of (i),(ii) and (iii), we prove the result (v). $\square$

Lemma 3. *For every fixed $p, q \in (0, 1)$, we have*

$$\frac{[n+m]_{p,q}[n+m-1]_{p,q}}{([n]_{p,q} + \beta)^2}q - 2\frac{[n+m]_{p,q}}{[n]_{p,q} + \beta} + 1 \leq \left(\frac{(p^n + q^n)[m]_{p,q} - \beta}{[n]_{p,q} + \beta} \right)^2.$$

Proof. Since $[n+m]_{p,q}[n+m-1]_{p,q} \leq [n+m]_{p,q}^2$, we get

$$\begin{aligned} \frac{[n+m]_{p,q}[n+m-1]_{p,q}}{([n]_{p,q} + \beta)^2}q - 2\frac{[n+m]_{p,q}}{[n]_{p,q} + \beta} + 1 &\leq \left(\frac{[n+m]_{p,q}}{[n]_{p,q} + \beta} - 1 \right)^2 \\ &= \frac{1}{([n]_{p,q} + \beta)^2} \left\{ \frac{p^{n+m} - q^{n+m}}{p-q} - \frac{p^n - q^n}{p-q} - \beta \right\}^2 \\ &= \frac{1}{([n]_{p,q} + \beta)^2} \left\{ \frac{(p^n + q^n)(p^m - q^m) - q^n(1 - p^m) - p^n(1 - q^m)}{p-q} - \beta \right\}^2 \\ &= \frac{1}{([n]_{p,q} + \beta)^2} \left\{ (p^n + q^n)[m]_{p,q} - \beta - \frac{q^n(1 - p^m) + p^n(1 - q^m)}{p-q} \right\}^2 \\ &\leq \frac{((p^n + q^n)[m]_{p,q} - \beta)^2}{([n]_{p,q} + \beta)^2} \quad \text{since, } 0 < q < p \leq 1. \end{aligned}$$

$\square$

Remark 1. *As a result of Lemma 2 and 3, we have*

$$\begin{aligned} C_{n,m}^{(\alpha,\beta)}((t-x)^2; x, p, q) &\leq \left(\frac{((p^n + q^n)[m]_{p,q} - \beta)^2}{([n]_{p,q} + \beta)^2} \right) x^2 + \left(\frac{[n+m]_{p,q}(2\alpha + p^{n-1})}{([n]_{p,q} + \beta)} - 2\alpha \right) \frac{b_n x}{([n]_{p,q} + \beta)} \\ &\quad + \frac{\alpha^2 b_n^2}{([n]_{p,q} + \beta)^2}. \end{aligned}$$

Lemma 4. *Taking supremum on $[0, b_n]$ on Second central moment, we get*

$$C_{n,m}^{(\alpha,\beta)}((t-x)^2; x, p, q) \leq b_n^2 \left\{ \frac{((p^n + q^n)[m]_{p,q} - \beta)^2}{([n]_{p,q} + \beta)^2} + \left| \frac{[n+m]_{p,q}(2\alpha + p^{n-1})}{([n]_{p,q} + \beta)^2} - \frac{2\alpha}{([n]_{p,q} + \beta)} \right| + \frac{\alpha^2}{([n]_{p,q} + \beta)^2} \right\}.$$

3. Korovkin-type approximation theorem

Assume C_ρ is the space of all continuous functions f such that

$$|f(x)| \leq M\rho(x), \quad -\infty < x < \infty.$$

Then C_ρ is a Banach space with the norm

$$\|f\|_\rho = \sup_{-\infty < x < \infty} \frac{|f(x)|}{\rho(x)}.$$

The subsequent results are used for proving Korovkin approximation theorem on unbounded sets.

Theorem 1. (See [2]) *There exists a sequence of positive linear operators U_n , acting from C_ρ to C_ρ , satisfying the conditions*

- (1) $\lim_{n \rightarrow \infty} \|U_n(1; x) - 1\|_\rho = 0$,
- (2) $\lim_{n \rightarrow \infty} \|U_n(\phi; x) - \phi\|_\rho = 0$,
- (3) $\lim_{n \rightarrow \infty} \|U_n(\phi^2; x) - \phi^2\|_\rho = 0$,

where $\phi(x)$ is a continuous and increasing function on $(-\infty, \infty)$ such that $\lim_{x \rightarrow \pm\infty} \phi(x) = \pm\infty$ and $\rho(x) = 1 + \phi^2$, and there exists a function $f^* \in C_\rho$ for which $\overline{\lim}_{n \rightarrow \infty} \|U_n f^* - f^*\|_\rho > 0$.

Theorem 2. (See [2]) *Conditions (1),(2),(3) of above theorem implies that*

$$\lim_{n \rightarrow \infty} \|U_n f - f\|_\rho = 0$$

for any function f belonging to the subset $C_\rho^0 := \{f \in C_\rho : \lim_{|x| \rightarrow \infty} \frac{|f(x)|}{\rho(x)} \text{ is finite} \}$.

Consider the weight function $\rho(x) = 1 + x^2$ and operators:

$$U_{n,m}^{\alpha,\beta}(f; x, p, q) = \begin{cases} C_{n,m}^{\alpha,\beta}(f; x, p, q) & \text{if } x \in [0, b_n], \\ f(x) & \text{if } x \in [0, \infty) \setminus [0, b_n]. \end{cases}$$

Thus for $f \in C_{1+x^2}$, we have

$$\begin{aligned} \|U_{n,m}^{\alpha,\beta}(f; \cdot, p, q)\| &\leq \sup_{x \in [0, b_n]} \frac{|U_{n,m}^{\alpha,\beta}(f; x, p, q)|}{1 + x^2} + \sup_{b_n < x < \infty} \frac{|f(x)|}{1 + x^2} \\ &\leq \|f\|_{1+x^2} \left[\sup_{x \in [0, \infty)} \frac{|U_{n,m}^{\alpha,\beta}(1 + t^2; x, p, q)|}{1 + x^2} + 1 \right]. \end{aligned}$$

Now, using Lemma 2 we will obtain,

$$\|U_{n,m}^{\alpha,\beta}(f; \cdot, p, q)\|_{1+x^2} \leq M \|f\|_{1+x^2}$$

if $0 < q < p \leq 1$, $\lim_{n \rightarrow \infty} p^n = \lim_{n \rightarrow \infty} q^n = N$, $N < \infty$ and $\lim_{n \rightarrow \infty} \frac{b_n}{[n]} = 0$.

Theorem 3. *For all $f \in C_{1+x^2}^0$, we have*

$$\lim_{n \rightarrow \infty} \|U_{n,m}^{\alpha,\beta}(f; \cdot, p_n, q_n) - f(\cdot)\|_{1+x^2} = 0$$

provided that $p := (p)_n$, $q := (q)_n$ with $0 < q_n < p_n \leq 1$, $\lim_{n \rightarrow \infty} p_n = 1$, $\lim_{n \rightarrow \infty} q_n = 1$, $\lim_{n \rightarrow \infty} p_n^n = \lim_{n \rightarrow \infty} q_n^n = N$, $N < \infty$, $\lim_{n \rightarrow \infty} [n]_{p_n, q_n} = \infty$ and $\lim_{n \rightarrow \infty} \frac{b_n}{[n]} = 0$.

Proof. Using the results of Theorem 1 and Lemma 2 (i),(ii) and (iii), we will achieve the following assessments, respectively:

$$\begin{aligned} \sup_{x \in [0, \infty)} \frac{|U_{n,m}^{\alpha,\beta}(1; x, p_n, q_n) - 1|}{1 + x^2} &= \sup_{0 \leq x \leq b_n} \frac{|C_{n,m}^{\alpha,\beta}(1; x, p_n, q_n) - 1|}{1 + x^2} = 0 \\ \sup_{x \in [0, \infty)} \frac{|U_{n,m}^{\alpha,\beta}(t; x, p_n, q_n) - t|}{1 + x^2} &= \sup_{0 \leq x \leq b_n} \frac{|C_{n,m}^{\alpha,\beta}(t; x, p_n, q_n) - x|}{1 + x^2} \leq \sup_{0 \leq x \leq b_n} \frac{\left| \frac{[n+m]_{p_n, q_n}}{[n]_{p_n, q_n} + \beta} - 1 \right| x + \frac{\alpha b_n}{[n]_{p_n, q_n} + \beta}}{1 + x^2} \\ &\leq \left| \frac{[n+m]_{p_n, q_n}}{[n]_{p_n, q_n} + \beta} - 1 \right| + \frac{\alpha b_n}{[n]_{p_n, q_n} + \beta} \rightarrow 0 \end{aligned}$$

and

$$\begin{aligned} \sup_{x \in [0, \infty)} \frac{|U_{n,m}^{\alpha,\beta}(t^2; x, p_n, q_n) - t^2|}{1 + x^2} &= \sup_{0 \leq x \leq b_n} \frac{|C_{n,m}^{\alpha,\beta}(t^2; x, p_n, q_n) - x^2|}{1 + x^2} \\ &\leq \sup_{0 \leq x \leq b_n} \frac{1}{1 + x^2} \left| \frac{(q_n[n+m]_{p_n, q_n}[n+m-1]_{p_n, q_n}x^2 + [n+m]_{p_n, q_n}(2\alpha + p_n^{n-1})b_nx + \alpha^2b_n^2)}{([n]_{p_n, q_n} + \beta)^2} - x^2 \right| \\ &\leq \left\{ \left| \frac{(q_n[n+m]_{p_n, q_n}[n+m-1]_{p_n, q_n}}{([n]_{p_n, q_n} + \beta)^2} - 1 \right| + \left| \frac{[n+m]_{p_n, q_n}(2\alpha + p_n^{n-1})}{([n]_{p_n, q_n} + \beta)^2} \right| + \frac{\alpha^2}{([n]_{p_n, q_n} + \beta)^2} \right\} \rightarrow 0 \end{aligned}$$

whenever $n \rightarrow \infty$, because we have $\lim_{n \rightarrow \infty} p_n = \lim_{n \rightarrow \infty} q_n = 1$ and $\lim_{n \rightarrow \infty} [n]_{p_n, q_n} = \infty$ as $n \rightarrow \infty$. $\square$

Theorem 4. Assuming c as a positive and real number independent of n and f as a continuous function which vanishes on $[c, \infty)$. Let $p := (p_n), q := (q_n)$ with $0 < q_n < p_n \leq 1, \lim_{n \rightarrow \infty} p_n = \lim_{n \rightarrow \infty} q_n = 1, \lim_{n \rightarrow \infty} p_n^n = \lim_{n \rightarrow \infty} q_n^n = N < \infty$ and $\lim_{n \rightarrow \infty} \frac{b_n^2}{[n]_{p, q}} = 0$. Then we have

$$\lim_{n \rightarrow \infty} \sup_{0 \leq x \leq b_n} |C_{n,m}^{\alpha,\beta}(f; x, p_n, q_n) - f(x)| = 0.$$

Proof. From the hypothesis on f , it is bounded i.e. $|f(x)| \leq M$ ($M > 0$). For any $\epsilon > 0$, we have

$$\left| f \left(\frac{p_n^{n-k}[k]_{p_n, q_n} + \alpha}{[n]_{p_n, q_n} + \beta} b_n \right) - f(x) \right| < \epsilon + \frac{2M}{\delta^2} \left(\frac{p_n^{n-k}[k]_{p_n, q_n} + \alpha}{[n]_{p_n, q_n} + \beta} b_n - x \right)^2,$$

where $x \in [0, b_n]$ and $\delta = \delta(\epsilon)$ are independent of n . Operating with the operator 2.1 on both sides, we can conclude by using Theorem 3 and Lemma 3,

$$\begin{aligned} \sup_{0 \leq x \leq b_n} |C_{n,m}^{\alpha,\beta}(f; x, p_n, q_n) - f(x)| &\leq \epsilon + \frac{2M}{\delta^2} b_n^2 \left\{ \left| 1 - 2 \frac{[n+m]_{p_n, q_n}}{([n]_{p_n, q_n} + \beta)} + \frac{q_n[n+m]_{p_n, q_n}[n+m-1]_{p_n, q_n}}{([n]_{p_n, q_n} + \beta)^2} \right| \right. \\ &\quad \left. + \left| \frac{(2\alpha + p_n^{n-1})[n+m]_{p_n, q_n}}{([n]_{p_n, q_n} + \beta)} - 2\alpha \right| \frac{1}{([n]_{p_n, q_n} + \beta)} + \frac{\alpha^2}{([n]_{p_n, q_n} + \beta)^2} \right\}. \end{aligned}$$

Since $\frac{b_n^2}{[n]_{p_n, q_n}} = 0$ as $n \rightarrow \infty$, we have the desired result. $\square$

4. Rate of Convergence

We will find the rate of convergence in terms of the Lipschitz class $Lip_M(\gamma)$, for $0 < \gamma \leq 1$. Assume that $C_B[0, \infty)$ denote the space of bounded continuous functions on $[0, \infty)$. A function $f \in C_B[0, \infty)$ belongs to $Lip_M(\gamma)$ if

$$|f(t) - f(x)| \leq M|t - x|^\gamma, \quad t, x \in [0, \infty)$$

is satisfied.

Theorem 5. *Let $f \in Lip_M(\gamma)$, then*

$$C_{n,m}^{\alpha,\beta}(f; x, p, q) \leq M(\lambda_{n,p,q}(x))^{\gamma/2},$$

where $\lambda_{n,p,q}(x) = C_{n,m}^{\alpha,\beta}((t - x)^2; x, p, q)$.

Proof. Since $f \in Lip_M(\gamma)$, and the operator $C_{n,m}^{\alpha,\beta}(f; x, p, q)$ is linear and monotone,

$$\begin{aligned} |C_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| &= \left| \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \begin{bmatrix} n+m \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n}\right)^{k(n+m-k-1)} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \right. \\ &\quad \left. f\left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n\right) - f(x) \right| \\ &\leq \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \begin{bmatrix} n+m \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n}\right)^{k(n+m-k-1)} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left| f\left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n\right) - f(x) \right| \\ &\leq M \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \begin{bmatrix} n+m \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n}\right)^{k(n+m-k-1)} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left| \left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n\right) - x \right|^\gamma. \end{aligned}$$

Using Remark (1) and Hölder's inequality with the values $p = \frac{2}{\gamma}$ and $q = \frac{2}{2-\gamma}$, we get

$$\begin{aligned} &|C_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \\ &\leq \frac{M}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\left\{ \begin{bmatrix} n+m \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n}\right)^{k(n+m-k-1)} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n\right) - x \right)^2 \right\}^{\frac{\gamma}{2}} \right. \\ &\quad \left. \times \left\{ \begin{bmatrix} n+m \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n}\right)^{k(n+m-k-1)} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \right\}^{\frac{2-\gamma}{2}} \right] \\ &\leq M \left[\left\{ \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \begin{bmatrix} n+m \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n}\right)^{k(n+m-k-1)} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n\right) - x \right)^2 \right\}^{\frac{\gamma}{2}} \right. \\ &\quad \left. \times \left\{ \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \begin{bmatrix} n+m \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n}\right)^{k(n+m-k-1)} \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \right\}^{\frac{2-\gamma}{2}} \right] \end{aligned}$$

$$\begin{aligned}
&= M \left[C_{n,m}^{\alpha,\beta}((t-x)^2; x, p, q) \right]^{\frac{\gamma}{2}} \\
&\leq M(\lambda_{n,p,q}(x))^{\frac{\gamma}{2}}.
\end{aligned}$$

□

In order to obtain rate of convergence in terms of modulus of continuity $\omega(f; \delta)$, we assume that for any $f \in C_B[0, \infty)$ and $x \geq 0$, modulus of continuity of f is given by

$$\omega(f; \delta) = \max_{\substack{|t-x| \leq \delta \\ t, x \in [0, \infty)}} |f(t) - f(x)|. \quad (4.1)$$

Thus it implies for any $\delta > 0$

$$|f(x) - f(y)| \leq \omega(f; \delta) \left(\frac{|x - y|}{\delta} + 1 \right). \quad (4.2)$$

Theorem 6. *If $f \in C_B[0, \infty)$, we have*

$$|C_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \leq 2\omega(f; \sqrt{\lambda_{n,p,q}(x)}),$$

where $\omega(f; \cdot)$ is modulus of continuity of f and $\lambda_{n,p,q}(x)$ be the same as in Theorem 5.

Proof. Using triangular inequality, we get

$$\begin{aligned}
|C_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| &= \left| \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \right. \\
&\quad \left. f \left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n \right) - f(x) \right| \\
&\leq \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left| f \left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n \right) - f(x) \right|.
\end{aligned}$$

Now using (4.2) and Hölder's inequality, we get

$$\begin{aligned}
&|C_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \\
&= \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{\left| \frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x \right|}{\delta} + 1 \right) \omega(f; \delta) \\
&\leq \omega(f; \delta) \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \\
&\quad + \frac{\omega(f; \delta)}{\delta} \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left| \frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x \right|
\end{aligned}$$

$$\begin{aligned}
&= \omega(f; \delta) + \frac{\omega(f; \delta)}{\delta} \left\{ \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \begin{bmatrix} n+m \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \right. \\
&\quad \left. \left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x \right)^2 \right\}^{\frac{1}{2}} \\
&= \omega(f; \delta) + \frac{\omega(f; \delta)}{\delta} \{ C_{n,m}^{\alpha,\beta}((t-x)^2; x, p, q) \}^{1/2}.
\end{aligned}$$

Now choosing $\delta = \lambda_{n,p,q}(x)$ as in Theorem 5, we have

$$|C_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \leq 2\omega(f; \sqrt{\lambda_{n,p,q}(x)}).$$

□

Next we calculate the rate of convergence in terms of the modulus of continuity of the derivative of function.

Theorem 7. *If $f(x)$ has a continuous bounded derivative $f'(x)$ and $\omega(f'; \delta)$ is the modulus of continuity of $f'(x)$ in $[0, A]$, then*

$$\begin{aligned}
&|f(x) - C_{n,m}^{\alpha,\beta}(f; x, p, q)| \leq \\
&M \left(\left| \frac{[n+m]_{p,q}}{[n]_{p,q} + \beta} - 1 \right| A + \frac{\alpha b_n}{[n]_{p,q} + \beta} \right) + 2(B_{n,p,q}(\alpha, \beta))^{1/2} \omega(f'; (B_{n,p,q}(\alpha, \beta))^{1/2}),
\end{aligned}$$

where M is a positive constant such that $|f'(x)| \leq M$.

Proof. Using Mean Value Theorem, we have

$$\begin{aligned}
f\left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n\right) - f(x) &= \left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x\right) f'(\xi) \\
&= \left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x\right) f'(x) + \left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x\right) (f'(\xi) - f'(x)),
\end{aligned}$$

where ξ is a point between x and $\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n$. By using the above identity, we get

$$\begin{aligned}
&C_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x) \\
&= f'(x) \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \begin{bmatrix} n+m \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x \right) \\
&\quad + \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \begin{bmatrix} n+m \\ k \end{bmatrix}_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left(\frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x \right) (f'(\xi) - f'(x)).
\end{aligned}$$

Hence,

$$|C_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)|$$

$$\begin{aligned}
&\leq |f'(x)| |C_{n,m}^{\alpha,\beta}((t-x); x, p, q)| \\
&\quad + \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left| \frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x \right| |f'(\xi) - f'(x)| \\
&\leq M \left(\left| \frac{[n+m]_{p,q}}{[n]_{p,q} + \beta} - 1 \right| A + \frac{\alpha b_n}{[n]_{p,q} + \beta} \right) \\
&\quad + \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left| \frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x \right| |f'(\xi) - f'(x)| \\
&\leq M \left(\left| \frac{[n+m]_{p,q}}{[n]_{p,q} + \beta} - 1 \right| A + \frac{\alpha b_n}{[n]_{p,q} + \beta} \right) \\
&\quad + \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left| \frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x \right| \\
&\quad \times \omega(f'; \delta) \left(\frac{\left| \frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x \right|}{\delta} + 1 \right),
\end{aligned}$$

since

$$|\xi - x| \leq \left| \frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x \right|.$$

Using it, we have

$$\begin{aligned}
&|C_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \\
&\leq M \left(\left| \frac{[n+m]_{p,q}}{[n]_{p,q} + \beta} - 1 \right| A + \frac{\alpha b_n}{[n]_{p,q} + \beta} \right) \\
&\quad + \omega(f'; \delta) \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left| \frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x \right| \\
&\quad + \frac{\omega(f'; \delta)}{\delta} \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left| \frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x \right|^2.
\end{aligned}$$

Now using Cauchy-Schwarz inequality for the second term, we obtain

$$\begin{aligned}
&|C_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \\
&\leq M \left(\left| \frac{[n+m]_{p,q}}{[n]_{p,q} + \beta} - 1 \right| A + \frac{\alpha b_n}{[n]_{p,q} + \beta} \right) \\
&\quad + \omega(f'; \delta) \left(\frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{n+m-k-1} (p^s - q^s \frac{x}{b_n}) \left| \frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x \right|^2 \right)^{1/2}
\end{aligned}$$

$$\begin{aligned}
& + \frac{\omega(f'; \delta)}{\delta} \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n+m} \left[\begin{matrix} n+m \\ k \end{matrix} \right]_{p,q} p^{\frac{k(k-1)}{2}} \left(\frac{x}{b_n} \right)^k \prod_{s=0}^{(n+m-k-1)} \left(p^s - q^s \frac{x}{b_n} \right) \left| \frac{p^{n-k} [k]_{p,q} + \alpha}{[n]_{p,q} + \beta} b_n - x \right|^2. \\
& = M \left(\left| \frac{[n+m]_{p,q}}{[n]_{p,q} + \beta} - 1 \right| A + \frac{\alpha b_n}{[n]_{p,q} + \beta} \right) \\
& \quad + \omega(f'; \delta) \sqrt{C_{n,m}^{\alpha,\beta}((t-x)^2; x, p, q)} + \frac{\omega(f'; \delta)}{\delta} C_{n,m}^{\alpha,\beta}((t-x)^2; x, p, q).
\end{aligned}$$

Using Remark (1), we see

$$\begin{aligned}
& \sup_{0 \leq x \leq A} C_{n,m}^{(\alpha,\beta)}((t-x)^2; x, p, q) \\
& \leq \sup_{0 \leq x \leq A} \left(\frac{((p^n + q^n)[m]_{p,q} - \beta)^2}{([n]_{p,q} + \beta)^2} \right) x^2 + \left(\frac{[n+m]_{p,q}(2\alpha + p^{n-1})}{([n]_{p,q} + \beta)} - 2\alpha \right) \frac{b_n x}{([n]_{p,q} + \beta)} \\
& \quad + \frac{\alpha^2 b_n^2}{([n]_{p,q} + \beta)^2} \\
& \leq \left(\frac{((p^n + q^n)[m]_{p,q} - \beta)^2}{([n]_{p,q} + \beta)^2} \right) A^2 + \left(\frac{[n+m]_{p,q}(2\alpha + p^{n-1})}{([n]_{p,q} + \beta)} - 2\alpha \right) \frac{b_n A}{([n]_{p,q} + \beta)} \\
& \quad + \frac{\alpha^2 b_n^2}{([n]_{p,q} + \beta)^2} \\
& := B_{n,p,q}(\alpha, \beta).
\end{aligned}$$

Thus,


$$|C_{n,m}^{\alpha,\beta}(f; x, p, q) - f(x)| \leq M \left(\left| \frac{[n+m]_{p,q}}{[n]_{p,q} + \beta} - 1 \right| A + \frac{\alpha b_n}{[n]_{p,q} + \beta} \right) + \omega(f'; \delta) \left[(B_{n,p,q}(\alpha, \beta))^{1/2} + \frac{1}{\delta} B_{n,p,q}(\alpha, \beta) \right].$$

Choosing, $\delta := (B_{n,p,q}(\alpha, \beta))^{1/2}$, we get the desired result. $\square$

Example:

Approximation of $f(x) = x^2$

Approximation of $x^2 \in C_{1+x^2}[0, \infty)$ function within $[0, 0.1]$, obtained for $n = 10, 8$ with $m = 1$ and $b_n = (\log n)^2$ is represented by the following figure.



It is clear from the figure that approximation is readily obtained for lower values of n using (p, q) variant of Bernstein-Stancu-Schurer-Chlodowsky polynomials in small dimension as it is required in quantum physics.

Conflict of Interest: The authors declare that they have no competing interests regarding the publication of this manuscript.

References

References

- [1] S.N. Bernstein, Démonstration du théorème de Weierstrass fondée sur le calcul de probabilités, *Comm. Soc. Math. Kharkow* (2), 13 (1912/1913), pp. 1-2.
- [2] A.D. Gadjiev, The convergence problem for a sequence of positive linear operators on unbounded sets and theorems analogous to that of P.P. Korovkin, *Dokl. Akad. Nauk SSSR*, 218 (5), 1001-1004. English translation in *Sov. Math. Dokl.* 15 (5), 1974, pp. 1433-1436.
- [3] E. Ibikli, Approximation by Bernstein-Chlodowsky polynomials, *Haceteepe Journal of Mathematics and Statistics*, Vol. 32 (2003), pp. 1-5.
- [4] E. Ibikli, On Stancu type generalization of Bernstein-Chlodowsky polynomials, *Mathematica*, 42 (65), 2000, pp. 37-43.
- [5] H. Karsh, V. Gupta, Some approximation properties of q -Chlodowsky operators, *Appl. Math. Comput.* 195 (2008), pp. 220-229.

- [6] A. Lupaş, A q -analogue of the Bernstein operator, *Seminar on Numerical and Statistical Calculus, University of Cluj-Napoca*, 9 (1987), pp. 85-92.
- [7] M. Mursaleen, K.J. Ansari, A. Khan, On (p, q) -analogue of Bernstein operators, *Appl. Math. Comput.*, 266 (2015), pp. 874-882.
- [8] M. Mursaleen, K.J. Ansari, A. Khan, On (p, q) -analogue of Bernstein operators (Revised) arXiv:1503.07404v2 [math.CA], (2015).
- [9] M. Mursaleen, K.J. Ansari, A. Khan, Some approximation results by (p, q) -analogue of Bernstein-Stancu operators, *Appl. Math. Comput.* 264 (2015), pp. 392-402.
- [10] M. Mursaleen, K.J. Ansari, A. Khan, Some approximation results by (p, q) -analogue of Bernstein-Stancu operators (Revised), arXiv:submit/1440017 [math.CA], (2015).
- [11] M. Mursaleen, Md. Nasiruzzaman, N. Ashirbayev, Some approximation results on Bernstein-Schurer operators defined by (p, q) -integers, *J. Inequal. Appl.*, 249 (2015).
- [12] V.N. Mishra, K. Khatri, L.N. Mishra, Deepmala, Inverse result in simultaneous approximation by Baskakov-Durrmeyer-Stancu operators, *J. Inequal. Appl.*, 586 (2013).
- [13] M. Mursaleen, F. Khan, A. Khan, Approximation properties for modified q -Bernstein-Kantorovich operators, *Num. Funct. Analysis and Optimization*, 36 (9), 2015, pp. 1178-1197.
- [14] M. Mursaleen, K.J. Ansari, A. Khan, Stability of some positive linear operators on compact disk, *Acta Mathematica Scientia*, 2015, 35 B (6), pp. 1-9.
- [15] M. Mursaleen, F. Khan, A. Khan, Approximation properties for Kings type modified q -Bernstein-Kantorovich operators, *Math. Meth. Appl. Sci.* 2015, DOI: 10.1002/mma.3454.
- [16] G.M. Phillips, On generalized Bernstein polynomials, *Numerical Analysis, World Scientific, River Edge*, vol. 98 (1996), pp. 263-269.
- [17] V. Sahai, S. Yadav, Representations of two parameter quantum algebras and (p, q) -special functions, *J. Math. Anal. Appl.*, 335 (2007), pp. 268-279 .